



Ruston Soil Profile

Surface layer: dark grayish brown fine sandy loam Subsurface layer: pale brown fine sandy loam

Subsoil - upper: red clay loam

Subsoil - middle: yellowish red fine sandy loam

Subsoil - lower: red sandy clay loam

Ruston soils are used mainly for woodland consisting of southern pine and some hardwoods with understories of shrubs or grasses. Some areas of understory vegetation are managed as range, which provides grazing for cattle, and some are managed for wildlife habitat. A small acreage is used for cotton, corn, soybeans, small grain, truck crops, or pasture. A considerable portion of the formerly cultivated acreage has been converted to pasture or southern pine woodland. These soils occur on 733,714 acres in Louisiana.

The Ruston series consists of very deep, well drained, moderately permeable soils that formed in loamy marine or stream deposits. These soils are low in fertility and within the root zone have moderately high levels of exchangeable aluminum that are potentially toxic to some agricultural crops but are ideal for the production of loblolly, slash, and longleaf pine. The soils have slight limitations for woodland use and management.



The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD). To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call 202-720-5964 (voice or TDD). USDA is an equal opportunity provider and employer.

